REFERENCES:

- Baohong Fang, Qiyao Ling,Weifeng Zhao, Yunli Ma, Pengli Bai, QiangWei, Haifeng Li, Changsheng Zhao (2009); Modification of polyethersulfone membrane by grafting bovine serum albumin on the surface of polyethersulfone/poly(acrylonitrile-co-acrylic acid) blended membrane; Journal of Membrane Science 329 ; 46–55.
- Chowdary KPR, Ramesh K.V.R.N.S (1994).,Improvement of dissolution rate and efficiency of nifedipine by solid dispersion in PVP-NCC and HPC-MCC. **Ind. J. Pharm Sci.**;56(3):95-99.(7)
- Chiou W.L, Riegelman S (1971), Pharmaceutical applications of solid dispersion systems. **J. Pharm. Sci**.; 60:1281.1302.
- Chowdary KPR, Rao S.S.(2000) Dissolution rate and formulation studies on Solid dispersion of itrconazole**. Ind. J. Pharm Sci**;62(6):471-474.
- Charumanee S, Okonoki S, Sirithunyalug J. (2004) Improvement of the Dissolution Rate of Piroxicam by Surface Solid Dispersion**. CMU. Journal**; Vol. 3(2):77-84.
- Duncan C.Q.M, Paul R.G, Kett V.L, Hopton M.L. (1999) The relevance of the amorphous state to pharmaceutical dosage forms: glassy drugs and freeze dried systems. **Int. J. Pharm**; 179:179–207.
- Dastmalchi S, Garjani A, Maleki N, Sheikhee G, Baghchevan V, Jafari-Azad P, Valizadeh H, Barzegar-Jalali M.(2005)Enhancing dissolution, serum concentrations and hypoglycemic effect of glibenclamide using solvent deposition technique. **J. Pharm. Sci**;8(2):175-181.
- Friedrich H, Fussnegger B, Kolter K, Bodmeier B. (2006)Dissolution rate improvement of poorly water-soluble drugs obtained by adsorbing solutions of drugs in hydrophilic solvents onto high surface area carriers. **Eur. J. Pharm. Biopharm**; 62(2):171–177.
- Francesca Maestrelli, Marcos Garcia-Fuentes, Paola Mura, Maria Jose´ Alonso; (2006) A new drug nanocarrier consisting of chitosan and hydoxypropylcyclodextrin; European Journal of Pharmaceutics and Biopharmaceutics, 63 ;79–86.
- Guidance for Industry, (2000) Waiver of In Vivo Bioavailability and Bioequivalence studies for Immediate Release Solid Oral Dosage Forms Based on a Biopharmaceutics Classification System. **FDA/CDER**.
- Gupta P, Kakumanu V.K, Bansal A.K. (2004) Stability and Solubility of Celecoxib–PVP Amorphous Dispersions: A Molecular Perspective. **Pharm. Res**; 21: 1762-1769.

- Ghebremeskel A.N, Vemavarapua C, Lodayaa M. (2007) Use of surfactants as plasticizers in preparing solid dispersions of poorly soluble API: Selection of polymer–surfactant combinations using solubility parameters and testing the processability. **Int. J. Pharm**; 328(2):119-129.
- Garima chawla arvind k. Bansal,(2008) Improved dissolution of a poorly water soluble drug in solid dispersions with polymeric and non-polymeric hydrophilic additives; *Acta Pharm. 58* ,257–274.
- J Thimmasetty, CVS Subrahmanyam, BA Vishwanath, and PR Sathesh Babu;,(*2009*) Solubility Parameter Estimation of Celecoxib by Current Methods; *Asian J. Research Chem. 2(2): April.-June.*
- Kakumanu V.K, Bansal A.K. (2002) Enthalpy Relaxation Studies of Celecoxib Amorphous Mixtures. **Pharm. Res**;19: 1873-1877.
- Kaushal A.M, Gupta P. Bansal A.K.(2004) Amorphous Drug Delivery Systems: Molecular Aspects, Design, and Performance Crit. **Rev. in Ther. Drug Carrier Syst**; 21:133–193.
- Kerc J, Srcic S. (1995) Thermal analysis of glassy pharmaceuticals. **Thermochim Acta**; 248:81-95.
- Kamal Dua, M.V. Ramana, U.V. Singh Sara, M. Himaja, Abhinav Agrawal, Vaibhav Garg and Kavita Pabreja,(2007) Investigation of Enhancement of Solubility of Norfloxacin β-Cyclodextrin in Presence of Acidic Solubilizing Additives, *Current Drug Delivery*, *4;*21-25.
- Lipinski C.A,(2000) Drug-like properties and the causes of poor solubility and poor permeability. **J. Pharmacol. Toxicol. Methods**;44:235-249.
- Leuner C, Dressman J.(2000) Improving drug solubility for oral delivery using solid dispersions. **Eur. J. Pharm**;50:47–60.
- Michael H, Stephen T, Cathy F.(2003) Part 1: "Oral Delivery of Poorly Soluble Drugs Pharmaceutical Manufacturing and Packing Sourcer". **Summer Samedan Ltd.**;03.
- Madhusudhan B, Rambhau B, Gudsoorkar V.R, Shete J.S, Apte S.S.(2002) Studies on Sulphamethoxazole Solid Dispersion and their Tablets. **Ind. J. Pharm. Sci**;64(30):233-238**.**
- Manish K.G., Yin-Chao T., David Goldman, and Robin H. Bogner.(2002) Hydrogen Bonding with Adsorbent during Storage Governs Drug Dissolution from Solid-Dispersion Granules **Pharm. Res**.; 19:1663-1672.

- Mohamed R. Berbera, keiji minagawaa, masahiro katoha,Takeshi morib, masami tanakac,(2008), Nanocomposites of 2-arylpropionic acid drugs based on Mg–Al layered double hydroxide for dissolution enhancement; **European journal of pharmaceutical sciences ,**3 5 ;354–360.
- Mani Upreti , Ken Strassburger , You L. Chen , Shaoxiong Wu and Indra Prakash;(2011), Solubility Enhancement of Steviol Glycosides and Characterization of Their Inclusion Complexes with Gamma-Cyclodextrin; ***Int. J. Mol. Sci.***, *12*, 7529-7553.
- Otsuka M, Katol F, Matsuda Y. (2002) Physicochemical Stability of Cimetidine Amorphous Forms Estimated by Isothermal Microcalorimetry. **AAPS PharmSciTech**; 3:1-13.
- Prentis R.A, Lis Y, Walker S.R.(1988) Pharmaceutical innovation by the seven UK-owned pharmaceutical companies. **Br J. Clin. Pharmacol.** 1964–1985;25:387–96.
- Patel D.M, Shah R.R, Jagani P.D.(2003) Studies to Enhance Dissolution of Piroxicam. **Ind. J. Pharm Sci**;65(3):264-267**.**
- Praveen Chaudhari, Pramodkumar Sharma, Nilesh Barhate, Parag Kulkarni and Chetan Mistry; (2007),Solubility enhancement of hydrophobic drugs using synergistically interacting cyclodextrins and cosolvent; **current science**, 1586 vol. 92, no. 11, 10 june.
- Preeta Anand a, Harish B. Nair b, Bokyung Sung a, Ajaikumar B. Kunnumakkara a, Vivek R. Yadav a, Rajeshwar R. Tekmal b, Bharat B. Aggarwal;(2009)) Design of curcumin-loaded PLGA nanoparticles formulation with enhanced cellular uptake, and increased bioactivity in vitro and superior bioavailability in vivo; **Biochemical Pharmacology** xxx ,xxx–xxx.
- Purwa Jain, Achhrish Goel, Shweta Sharma, Meghal Parmar,(2010), Solubility Enhancement Techniques With Special Emphasis On Hydrotrophy; **International Journal Of Pharma Professional's** ,Issue 1, July**.**
- Ratul K.D, Naresh K, Utpal B;(2010), Encapsulation of curcumin in alginate-chitosan-pluronic composite nanoparticles for delivery to cancer cells**.** **Nanomedicine: Nanotechnology, Biology, and Medicine,** 6 ;153–160.
- Roul LK *, Manna NK , Parhi RN , Sahoo S and Suresh P;(2012), Dissolution rate enhancement of Alprazolam by Solid Dispersion; **Ind J Pharm Edu Res**, Jan-Mar, Vol 46/ Issue 1.
- Serajjuddin A.T.M, (1999) Solid dispersion of poorly water-soluble drugs:early promises, subsequent problems, and recent breakthrough. **J. Pharm.Sci**.88:1058-1066.

- Singla AK, Nagra L. (2000) Release enhancement and reduction in ulcerogenicity of naproxen by polyvinylpyrrolidone. **Ind. J. Pharm Sci**; 62(2) : 126-128.
- Shawn A. Mitchell, Thomas D. Reynolds, Tina P. Dasbach; (2003) A compaction process to enhance dissolution of poorly watersoluble drugs using hydroxypropyl methylcellulose; **International Journal of Pharmaceutics** 250; 3_11.
- Shirwaikar AA, Ramesh A. (2004) Fast disintegrating tablets of Atenolol by dry granulation method. **Ind. J. Pharm. Sci**. 66(4):422-428.
- Sharma S, Sher S, Badve S, Pawar A.P.(2005), Adsorption of Meloxicam on Porous Calcium Silicate: Characterization and Tablet Formulation. **AAPS PharmSciTech**:1-28.
- Shamkant L. Shimpi, Chauhan B, Mahadik K.R, Paradkar A. (2005) Stabilization and Improved in Vivo Performance of Amorphous Etoricoxib using Gelucire 50/13. **Pharm. Res**; 10: 1727-1734.
- Tang, Michael J, Pikal, Lynne S. Charlie T.X. (2002) A Spectroscopic Investigation of Hydrogen Bond Patterns in Crystalline and Amorphous Phases in Dihydropyridine Calcium Channel Blockers. **Pharm. Res**;19: 477-483.
- Tejal J. Shah, Avani F. Amin, Jolly R. Parikh,and Rajesh H. Parikh; (2007) Process Optimization and Characterization of Poloxamer Solid Dispersions of a Poorly Water-soluble Drug, **AAPS PharmSciTech** ; 8 (2) Article 29.
- Urbanetz N. A. (2006) Stabilization of solid dispersions of nimodipine and polyethylene glycol. **Eur. J. Pharm Sci.**;8:67-76.
- Vasanthavada M, Tong W.Q, Joshi Y, Kislaliogluc M.S.(2004), Phase Behavior of Amorphous Molecular Dispersions I: Determination of the Degree and Mechanism of Solid Solubility. **Pharm. Res**;21: 1598-1606.
- Yu L. X, Amidon G.L, Polli J. E, Zhao H, Mehta M, Conner D. P. Shah V. P, Lesko L. J, Chen M.L, Lee V. H. L and Hussain A. S.(2002) Biopharmaceutics Classification System: the scientific basis for biowaiver extension. **Pharm**. **Res**;19:921–925.
- Yellela S.R. Krishnaiah;(2010)Pharmaceutical Technologies for Enhancing Oral Bioavailability of Poorly Soluble Drugs; **Journal of Bioequivalence & Bioavailability** Volume 2(2): 028-036–028.

- Zhao N, Augsburger L. (2005) The influence of swelling capacity of superdisintegrant in different pH media on the dissolution of hydrochlorthiazide from directly compressed tablets. **AAPS Pharma.SciTech**;6(1):E120-E126.